



Stevens, E., Massey, R., Laabei, M., Gardner, S., & Somerville, G. (2017). Cytolytic toxin production by *Staphylococcus aureus* is dependent upon the activity of the protoheme IX farnesyltransferase. *Scientific Reports*, 7, [13744]. <https://doi.org/10.1038/s41598-017-14110-8>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1038/s41598-017-14110-8](https://doi.org/10.1038/s41598-017-14110-8)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via scientific reports at <https://www.nature.com/articles/s41598-017-14110-8>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

SCIENTIFIC REPORTS

OPEN

Cytolytic toxin production by *Staphylococcus aureus* is dependent upon the activity of the protoheme IX farnesyltransferase

Emily Stevens^{1,4}, Maisem Laabei^{1,2}, Stewart Gardner³, Greg A. Somerville³ & Ruth C. Massey⁴

Staphylococcus aureus is a medically important pathogen with an abundance of virulence factors that are necessary for survival within a host, including the production of cytolytic toxins. The regulation of toxin production is mediated by the Agr quorum sensing system, and a poorly defined post-exponential growth phase signal independent of Agr. As part of a recent genome wide association study (GWAS) to identify novel loci that alter the expression of cytolytic toxins, a polymorphism in the *cyoE* gene, which encodes a protoheme IX farnesyltransferase, was identified. This enzyme is essential for processing heme into the electron transport chain for use as an electron acceptor. Interestingly, without this enzyme *S. aureus* were repressed in their ability to secrete cytolytic toxins, and this appears to be mediated through repression of the Agr quorum sensing system. We hypothesize that the loss of electron transport is inducing feedback inhibition of metabolic capabilities that suppress the TCA cycle, and that this coupled with decreased RNAPIII transcription prevents synthesis of cytolytic toxins.

Staphylococcus aureus is a commensal bacterium that persistently colonises the nasal passages of approximately 20% of the human population^{1,2}. As an opportunistic pathogen, it commonly infects people whose immune system is compromised through illness, injury or age. These infections vary widely in terms of anatomic site and disease severity, ranging from minor skin and soft tissue infections to life threatening pneumonia or bacteraemia^{1,2}. Central to *S. aureus*' ability to survive the host immune response is the synthesis of numerous virulence determinants that help facilitate nutrient acquisition and immune evasion. Specifically, *S. aureus* produces adhesins that allow it to adhere to and colonise host tissues; proteins and a capsule that facilitate evasion of the host immune system; and secreted toxins that damage host cells and release nutrients. The expression of these virulence determinants is in part regulated by the Agr quorum sensing system, with a poorly defined post-exponential growth phase signal independent of Agr³. The tricarboxylic acid (TCA) cycle activity⁴ is also critical for cytolytic activity where TCA cycle mutants have decreased synthesis of secreted toxins. In staphylococci, TCA cycle activity is induced during the post-exponential growth phase, providing the bacterium with carbon, energy, and reduced dinucleotides⁵. TCA cycle-derived four- and five-carbon intermediates are used in biosynthetic reactions to synthesize precursors (e.g. amino acids); while energy (e.g. ATP) drives many critical cellular processes and reduced dinucleotides can donate electrons to the electron transport chain to generate ATP by oxidative phosphorylation.

Recently we analysed genetic polymorphisms within *S. aureus* clinical isolates from the major hospital-associated MRSA lineage ST239⁶, and found there was little variability in their adhesive capabilities, but there was significant variation in cytolytic toxin production⁷. The application of GWAS (genome wide association studies) to this data identified a number of polymorphic loci that were statistically associated with altering cytolytic toxin synthesis. One of those loci, *cyoE*, encodes the enzyme protoheme IX farnesyltransferase, an enzyme involved in catalysing the conversion of heme B to heme O^{8,9}. Heme O is incorporated into the electron transport chain as an electron acceptor, facilitating aerobic respiration and energy production^{10,11}. As mentioned, TCA

¹Milner Centre for Evolution, Dept. of Biology and Biochemistry, University of Bath, Bath, UK. ²Microbiology Department, Hospital Universitari Germans Trias i Pujol, Institut d'Investigació Germans Trias i Pujol, Universitat Autònoma de Barcelona, Badalona, Spain. ³School of Veterinary Medicine and Biomedical Sciences, University of Nebraska-Lincoln, Nebraska, USA. ⁴School of Molecular and Cellular Medicine, University of Bristol, Bristol, UK. Correspondence and requests for materials should be addressed to R.C.M. (email: ruth.massey@bristol.ac.uk)




Figure 1. GWAS identified association between mutations in the *cyoE* gene and toxicity. **(A)** Result of a GWAS study on a collection of 90 ST239 MRSA isolates that associated polymorphisms in the *cyoE* gene with the toxicity of the bacteria. The data presented is the mean toxicity (as measured by lysis of fluorescent dye containing vesicles²² of the isolates with and without polymorphisms in the gene. Eight isolates contained the TW20 reference *cyoE* gene, and 82 isolates contained the SNP. **(B)** The position of the amino acid change P87L in the protein, protoheme IX farnesyltransferase, encoded by the *cyoE* gene has been mapped to a ribbon model of the structure of this enzyme, and in **(C)** colorimetric surface colouring has been used to illustrate the white hydrophobic domains of the protein, the blue positively charged amino acid residues, and the red negatively charged amino acid residues. **(D)** A model of the protoheme IX farnesyltransferase situated in the cell membrane, and position of the toxicity associated amino acid change, as modelled by Protter¹⁶.




Figure 2. Functional verification of the contribution the *cyoE* gene makes to *S. aureus* toxicity. The *cyoE* gene was inactivated by transposon insertion in both the JE2 and SH1000 backgrounds. In the JE2 background the loss of toxicity was restored by expressing the gene from an inducible promoter on the pRMC2 vector plasmid. Complementation controls of the empty vector (JE2 Δ cyoE (pRMC2)), and the vector with the *cyoE* gene cloned, but uninduced (JE2 Δ cyoE (pCyoE)) have been included.

cycle mutants have decreased accumulation of cytolytic toxins; hence, the connection between the electron transport chain and the TCA cycle suggests these variants may have decreased TCA cycle activity and toxin accumulation. To address these possibilities, the metabolism and phenotype of a *cyoE*-deficient mutant and complemented strains were analysed.

Materials and Methods

Strains and cultivation conditions for toxicity assays. The Δ *cyoE* mutant (NE1434) was obtained from the Nebraska Transposon Mutant Library, which is a collection of 1,952 *S. aureus* mutants in the USA300 strain JE2¹². The USA300 JE2 wild type was used as a control in all assays. Bacterial strains were grown at 37 °C in brain heart infusion broth (BHI; Oxoid) or on tryptic soy agar (TSA; Sigma). When needed, erythromycin (5 µg/ml), chloramphenicol (10 µg/ml) and tetracycline (50 ng/ml) were added.

Growth, pH and acetic acid assays. Unless stated otherwise, bacterial strains were grown in filter-sterilized tryptic soy broth (TSB; Becton Dickinson and Company) and cultivated at 37 °C, with 225 rpm aeration, and using a flask-to-medium ratio of 10:1. Bacterial pre-cultures were prepared from overnight cultures diluted 1:100 in TSB and incubated for 1.5 to 2 h. These pre-cultures were centrifuged for 5 min at 5,000 rpm, and the exponentially growing cells were inoculated into pre-warmed TSB to an optical density at 600 nm (OD₆₀₀) of 0.02. No antibiotics were used in this assay. Cultures were diluted prior to reading the density at the later stages of growth to avoid any saturation effects in the spectrophotometer. The pH of the culture medium was determined hourly using an Accumet AR60 pH meter (Fisher Scientific). The acetic acid assays were performed on culture supernatants (1 ml) that were harvested hourly by centrifugation and the acetate and glucose concentrations were determined with kits purchased from R-Biopharm and used according to the manufacturer's protocol.

Toxicity assay. THP-1 cells are an immortal monocytic cell line¹³ that is sensitive to 13 of the 15 toxins produced by *S. aureus* that are present in the bacterial supernatant⁷. They are continuously sub-cultured at 2–3 day intervals in a solution of RPMI 1640 containing fetal bovine serum and an antibiotic solution of 200 mM L-glutamine, 10,000 units of penicillin and 10 mg/ml streptomycin. Following overnight growth in BHI broth, the bacterial cultures were centrifuged for 10 minutes at 10,000–12,000 × g and the supernatant was harvested. The supernatant was diluted to a 30% vol/vol in BHI broth and 20 µl of this was added to 20 µl of washed THP-1 cells at a concentration of 120–150 cells per 1 µl, and incubated for 12 minutes at 37 °C. Following incubation of bacterial supernatant with THP-1 cells, samples were stained with 260 µl Guava ViaCount reagent, incubated at room temperature for 5 minutes, and loaded onto a Guava flow cytometer to determine the percentage of THP-1 cell death in each sample.

Complementation of Tn mutant. To confirm that loss of the *cyoE* gene was responsible for the observed loss in toxicity, the wild-type *cyoE* gene was re-introduced into the transposon mutant. The plasmid vector pRMC2 was used because it contains a tetracycline-inducible promoter region that allows transcription of the gene of interest to be controlled. The wild-type *cyoE* gene was amplified by PCR using the following primer sequences:

cyoE FW: GCTGGTACCATGAACAAATTTAAGGAG; *cyoE* RV: GCGAATTCATTTTCATCCTAACTTAATT

Restriction enzyme sites for KpnI and EcoRI were added to the forward and reverse primers, respectively. The *cyoE* gene and plasmid pRMC2 were then digested with KpnI and EcoRI and the resultant products were ligated using T4 DNA Ligase. Successfully ligated plasmids containing the wild-type *cyoE* gene were transformed into *E. coli* DH5 α competent cells through electroporation, plasmid DNA was isolated and passed through *S. aureus* RN4220, before finally being transformed into the strain JE2 Δ *cyoE* transposon mutant NE1434. For




Figure 3. Activation of the Agr quorum sensing system is dependent upon protoheme IX farnesyltransferase activity. **(A)** The growth of strains carrying the RNIII reporter plasmid was monitored over 12 hours demonstrating that inactivation of the *cyoE* gene has minimal effect on the growth of *S. aureus* during this time frame *in vitro*. **(B)** Expression of Agr was monitored over 12 hours of growth where the inactivation of *cyoE* was found to have a negative effect on Agr activity. Control strains RN6390B (Agr+) and RN6911 (Agr−) have been included for comparison. **(C)** The downstream effect of the loss of Agr activity was verified by comparing the production of alpha toxin and the PSMs by both the wild-type and Δ *cyoE* mutant, where the mutant produced 2.2-fold less alpha toxin and 1.5-fold less PSM.

transformation through electroporation, bacteria were cultivated in BHI liquid culture to an OD₅₅₀ of 0.2–0.3 and washed four times in ice cold 0.5 M sucrose. After the final wash bacteria were suspended in 100 μ l of 0.5 M sucrose before being added to 1–5 μ g/ml of DNA. Bacteria were incubated on ice with the DNA for 20 minutes and electroporated in 0.2 cm cuvettes for 4.2–4.6 milliseconds. Following electroporation, 800 μ l BHI was added to the cuvettes and incubated for 1 hour at 37 °C without shaking. The transformants were then plated on TSA containing 10 μ g/ml chloramphenicol, and for the transposon mutant strain 5 μ g/ml erythromycin was also added to the agar.

Phage transduction of Δ *cyoE* from strain JE2 into strain SH1000. Donor cells were inoculated into liquid culture from single colonies and grown overnight, and the following day 200 μ l of this culture was added to 25 ml BHI containing 250 μ l 1 M MgSO₄ and 250 μ l 1 M CaCl₂. This was grown for one hour and then 100 μ l phage 11 was added to the culture and grown for a further four hours minimum. Supernatant was obtained from this culture through centrifugation (12,000 \times g for 3 minutes), and was then filter sterilised. Optimal plaque titre was in the range of 10^{7–10}. Next, recipient cells were grown overnight in 20 ml LK broth (1% tryptone, 0.5% yeast extract, 0.7% potassium chloride), then this culture was centrifuged (2,500 \times g for 10 minutes) and the pellet suspended in 1 ml LK broth. To 250 μ l of recipient cells was added 500 μ l LK broth plus 10 mM CaCl₂ and 250 μ l of the phage lysate from the previous step. This culture was incubated statically for 25 minutes and then with shaking at 180 rpm for 15 minutes. 500 μ l ice cold 0.02 M sodium citrate was then added, and the culture centrifuged at 10,000 \times g for 10 minutes. The pellet was suspended in 500 μ l 0.02 M sodium citrate and left on ice for 2 hours.




Figure 4. The activity of the TCA cycle is affected by the loss of expression of protoheme IX farnesyltransferase. The pH (A) and acetate (B) levels of the supernatant of the wild type (JE2), $\Delta cyoE$ mutant and the complemented mutant were quantified over 12 hours of growth. Inactivation of the *cyoE* gene affected both the pH and acetate levels, the effects of which were complemented by expression of the gene from the plasmid pCyoE. (C) Aconitase activity, a key feature of the TCA cycle, was quantified in the wild type (JE2), $\Delta cyoE$ mutant and complemented mutant. Inactivation of the *cyoE* gene significantly reduced aconitase activity, the effect of which was complemented by expression of the gene from the pCyoE plasmid.

100 μ l of this was then plated neat on LKA plates containing 0.02 M sodium citrate and selective antibiotic, and this was incubated for at least 20 hours at 37 °C.

RNAIII activity assay. A plasmid containing a GFP-tagged copy of the RNAIII gene was transformed into strains USA300 JE2, $\Delta cyoE$, RN6390B (an *agr*-positive strain) and RN6911 (an *agr*-negative strain). Single colonies were then inoculated into BHI as described above for liquid overnight cultures, and the following morning a 1:10,000 dilution was made into BHI at a flask to medium ratio of 10:1. Strains were cultured at 37 °C and aerated at 180 rpm; OD₆₀₀ and GFP fluorescence (485/520) readings were then taken at hourly intervals over 12 hours.




Figure 5. Repression of the TCA cycle by growing *S. aureus* under microaerobic conditions repressed the expression of cytolytic toxins. The wild type (JE2) strain was grown under microaerobic (10:8 flask to broth volume ratio) and under aerobic (10:1 flask to broth volume ratio) to determine the effect this has on TCA cycle activity has on cytolytic toxin activity. An Agr mutant and the Δ cyoE mutant both grown aerobically were included. Growth of the wild type strain under the microaerobic conditions had an equivalent effect on toxicity as inactivation of either the Agr system or the cyoE gene, demonstrating the contribution the TCA cycle makes to the toxicity of *S. aureus*.

Aconitase activity. Bacteria were harvested during the postexponential growth phase (6 h) by centrifugation, suspended in ACN buffer (100 mM fluorocitrate, 90 mM Tris/HCl, pH 8.0), and lysed with lysing matrix B tubes and a FastPrep instrument (MP Biomedicals). The lysate was centrifuged for 5 min at 13,200 rpm at 4 °C, and the aconitase activity in the cell-free lysate was measured by the method of Kennedy *et al.*¹⁴. One unit of aconitase activity is defined as the amount of enzyme necessary to give a $A_{240} \text{ min}^{-1}$ of 0.0033.

Western blot for α -toxin. Proteins were precipitated from bacterial supernatant following 18 hrs of growth using trichloroacetic acid (TCA) at a final concentration of 20% for 1 hour on ice. Samples were then washed three times using ice cold acetone, and solubilised in 100 μ l 8 M urea. 20 μ l of each sample was mixed with 20 μ l loading dye, and heated at 100 °C for 2 minutes. 10 μ l of each sample was then subjected to 10% SDS-PAGE and separated proteins were electroblotted onto a nitrocellulose membrane using a semi-dry blotter at 15 V for 30 minutes (BioRad). Membranes were blocked overnight using 3% BSA in PBS-T (containing 0.1% Tween), and were then incubated for 1 hour with rabbit polyclonal antibodies specific for α -toxin. After washing 3 times for 5 minutes with PBS, membranes were incubated for another hour with horseradish peroxidase-coupled Protein G. All incubation steps were done at room temperature. Membranes were washed twice for 20 minutes in PBS, and blots were then visualised using an Opti-4CN detection kit. Band intensities were quantified using ImageJ (v 1.46r).

PSM quantification. Overnight cultures were diluted 1:1000 in 50 ml BHI and grown for 18 hours at 37 °C with shaking (180 rpm). 30 ml supernatant was added to 10 ml 1-butanol and these samples were incubated for 3 hours at 37 °C with shaking. Samples were then centrifuged for 3 minutes and 1 ml of the upper organic phase was collected. Protein samples were concentrated overnight using a SpeedVac and dried samples were then solubilised in 150 μ l 8 M urea. Samples were loaded and run on 10% SDS-PAGE as described above and then stained using SimplyBlue SafeStain as per the protocol. Band intensities were quantified using ImageJ (v 1.46r).

Micro-aerobic environment. To assess the effect of microaerobic growth conditions on toxicity in the wild-type strain JE2, bacteria were cultivated using a flask-to-medium ratio of 10:8 and toxicity assays were conducted using culture supernatant as described above. All other growth conditions remained unchanged.

Statistics. All of the data presented here was found to be normally distributed, and as such significance (p values) were determined using the Student's unpaired 2-tailed T-test.

Results and Discussion

Association between toxicity and the polymorphic cyoE gene. A recent GWAS study identified an association between a polymorphic version of the cyoE gene with changes in the cytolytic activity (toxicity) of *S. aureus*, suggesting this locus may contribute to toxicity⁷. A comparison of these data has demonstrated that there was a 2.9-fold reduction (Fig. 1A; $p = 0.0008$) in the mean toxicity of *S. aureus* containing the SNP in the cyoE gene relative to those with the cyoE gene with no SNPs (i.e. that found in the reference strain of the ST239 lineage, TW20⁶). The cyoE gene encodes a protoheme IX farnesyltransferase^{8,9}, which is a membrane associated protein involved in the processing of heme, enabling the bacteria to respire aerobically^{10,11}.

The SNP change observed in the collection of clinical strains confers a change from proline to leucine at position 87 in the translated product of the *cyoE* gene. To determine the likelihood of this change affecting the activity of this protein we built a model of it using SWISS-MODEL¹⁵. We then viewed this model in Chimera to visualise where the amino acid change occurred and how it might therefore affect the activity of the protein in the clinical strains (Fig. 1B and C). As this is a membrane protein we also present a model of this which was generated using Protter¹⁶ (Fig. 1D). Based on these we hypothesise that the change to leucine could give the loop in which it is located more flexibility and make it more hydrophobic, which could result in the loop flipping inwards on itself. The change also appears to be within a region that could be the heme binding site of this enzyme; Fig. 1C shows the suggested structure of this protein with coulombic surface colouring to show positively and negatively charged regions of the protein. The blue region in the lower middle of the structure is thought to be the active site of this enzyme, thus any change to the structure of the loop beneath, particularly the loop flipping inwards, could affect the function of the active site. Further analysis of the structure of this protein would be required to confirm this hypothesis.

Functional verification of the contribution protoheme IX farnesyltransferase makes to *S. aureus* toxicity. To verify the association between the *cyoE* gene and toxicity, the cytolytic activity of a Δ *cyoE* transposon mutant from the Nebraska Transposon Mutant library and the isogenic strain JE2 were compared. To quantify this, the THP-1 monocytic cell line, which is sensitive to both the Phenol Soluble Modulins (PSMs) and many of the other cytolytic toxins secreted by *S. aureus*, was exposed to culture supernatants and toxicity was assessed. As suggested by the GWAS results, the Δ *cyoE* mutant had significantly decreased toxicity relative to the isogenic wild-type strain (Fig. 2; $p < 0.0001$). Complementation of the Δ *cyoE* mutant resulted in restoration of toxicity to wild type levels. To ascertain if the effect of inactivating *cyoE* was strain-dependent, the mutation was transduced into *S. aureus* strain SH1000 and toxicity was assessed. Similar to the strain JE2 background, inactivation of the *cyoE* gene in strain SH1000 caused a significant loss of toxicity (Fig. 2; $p = 0.004$). Taken together, these data confirm that the *cyoE* gene contributes to the ability of *S. aureus* to produce toxins.

Protoheme IX farnesyltransferase activity affects the ability to activate the Agr quorum sensing system. As the Agr quorum sensing system is a major regulator of toxin synthesis^{17–19} and as the inactivation of *cyoE* dramatically affects toxicity, we hypothesised that this effect of the loss of *cyoE* may be mediated through the Agr system. To test this hypothesis, an RNAIII::gfp fusion plasmid, which acts as a reporter of Agr activity, was introduced into the JE2 wild-type and Δ *cyoE* mutant strains and fluorescence was monitored over time (Fig. 3A and B). There was a significant reduction in fluorescence in the Δ *cyoE* mutant ($p = 0.025$), demonstrating that RNAIII transcription and consequently Agr activation is altered in the Δ *cyoE* mutant relative to the wild-type strain. To further examine the effect of the loss of the *cyoE* gene on Agr activity we quantified the expression of toxins known to be under its regulation, where the secretion of alpha toxin was quantified by western blotting and PSMs by butanol extraction. These assays were performed in triplicate (Fig. 3C) where we found on average there was 2.2-fold more alpha toxin, and 1.5-fold more PSMs expressed by the wild type JE2 strains when compared to the Δ *cyoE* mutant. These results confirm that in the absence of the *cyoE* gene the expression and activity of the Agr quorum sensing system is repressed.

Protoheme IX farnesyltransferase affects the TCA cycle. The protoheme IX farnesyltransferase (CyoE) is essential for electron transport in many organisms, and electron transport facilitates the oxidation of reduced dinucleotides that are generated from TCA cycle activity. As such the inactivation of *cyoE* in *S. aureus* should dramatically decrease TCA cycle activity. The decreased Agr activity (Fig. 3) coupled with decreased TCA cycle activity could provide a potential explanation for the decreased cytotoxicity we observed for the Δ *cyoE* mutant, as we have previously shown secretion of cytotoxins requires TCA cycle activity⁴. To test this hypothesis, the catabolism of acetate, which requires TCA cycle activity, was assessed. Acetate accumulates in the culture medium during the exponential growth phase due to the incomplete oxidation of carbohydrates when the TCA cycle is repressed. The catabolism of acetate begins when carbohydrates are depleted and the TCA cycle is de-repressed in the post-exponential growth phase. Loss of a functional TCA cycle will result in a deficiency in acetate catabolism, leading to a build-up of acetate in the culture medium and lowering its pH. As expected, the acidification of the media was equivalent, as was the accumulation of acetate between the wild-type, Δ *cyoE* mutant and complemented strains over the first five hours of growth (Fig. 4A). However, once the bacteria ceased growing exponentially (from 5hr onwards) the Δ *cyoE* mutant failed to alkalise the culture medium and did not catabolise acetate (Fig. 4B). To specifically verify the effect of *cyoE* on the TCA cycle activity we also quantified aconitase activity, which is a major TCA cycle enzyme catalysing the interconversion of citrate and isocitrate. During the post-exponential growth phase (6hr) we found that the wild-type and complemented mutant strains both had significantly more aconitase activity when compared to the Δ *cyoE* mutant (Fig. 4C; $p = 0.001$). Taken together, the Δ *cyoE* mutant's inability to catabolise acetate and alkalise the culture medium, demonstrates that inactivation of *cyoE* blocks TCA cycle activity.

Growth of *S. aureus* under micro-aerobic conditions mimics the effect of a loss of *cyoE*. As an alternative means of demonstrating the effect of reducing TCA cycle activity on toxicity, we examined whether an effect equivalent to the inactivation of *cyoE* could be achieved by culturing the bacteria in a microaerobic environment. As a facultative anaerobic bacterium, *S. aureus* can grow under anaerobic conditions, however its growth rate is significantly affected with one study finding a nine-fold difference in final cell density when aerobic and anaerobic growth conditions were compared²⁰. This study also found that under anaerobic conditions *S. aureus* secretes less alpha toxin, however these effects on cell growth make assessing the relative levels of expression of quorum-sensing dependent proteins complicated. As such, to repress the TCA cycle while minimising

the growth defects associated with anaerobic conditions we created a micro-aerobic environment by growing the bacteria in air, but manipulating the flask-to-medium ratio²¹. We grew JE2 for 18 hrs and the effect of comparing a flask-to-medium ratio of 10:1 and 10:8 on the secretion of toxins was quantified (Fig. 5). The bacteria grown in micro-aerobic conditions had a relatively small (1.9-fold) decrease in biomass when compared to those grown aerobically. There was however a significant effect on toxicity with the bacteria growing in a micro-aerobic environment expressing levels of toxicity equivalent to an aerobically grown $\Delta agrB$ mutant (Fig. 5).

Conclusion

S. aureus is one of the most sequenced bacterial pathogens, with many thousands of genomes being publicly available. We are at a point in which we can utilise this sequence data to understand the biology of this important pathogen in greater detail. This study began as a GWAS analysis of a collection of ST239 isolates and is completed here as we characterise the molecular detail of the role of a GWAS identified locus on virulence. Here we demonstrate that the *cyoE* gene, which encodes a protoheme IX farnesyltransferase enzyme, plays a critical role in the ability of *S. aureus* to secrete cytolytic toxins. In the $\Delta cyoE$ mutant, activation of the Agr quorum sensing system is significantly delayed, despite the bacteria reaching sufficient cell density. We believe this effect on the Agr system is coupled to repression of the TCA cycle caused by the loss of heme O production and its role as an electron acceptor in the electron transport chain. This work reiterates the important link between metabolism and virulence in *S. aureus*, but also demonstrates the variability that exists in these attributes amongst clinical isolates causing disease in humans. As genome sequencing becomes more embedded in clinical diagnostic procedures, information relating to such polymorphic loci could be used to assist in the diagnosis of highly virulent infections.

References

- Gordon, R. J. & Lowy, F. D. Pathogenesis of methicillin-resistant *Staphylococcus aureus* infection. *Clin. Infect. Dis.* **46**, 350–359 (2008).
- Naber, C. K. *Staphylococcus aureus* bacteremia: Epidemiology, pathophysiology, and management strategies. *Clin. Infect. Dis.* **48**, 231–237 (2009).
- Vandenesch, F., Kornblum, J. & Novick, R. P. A temporal signal, independent of *agr*, is required for *hla* but not *spa* transcription in *Staphylococcus aureus*. *J. Bacteriol.* **173**(20), 6313–6320 (1991).
- Somerville, G. A. *et al.* *Staphylococcus aureus* aconitase inactivation unexpectedly inhibits post-exponential-phase growth and enhances stationary-phase survival. *Infect. Immun.* **70**(11), 6373–6382 (2002).
- Somerville, G. A. *et al.* Correlation of acetate catabolism and growth yield in *Staphylococcus aureus*: implications for host-pathogen interactions. *Infect. Immun.* **71**(8), 4724–4732 (2003).
- Holden, M. T. G. *et al.* Genome sequence of a recently emerged, highly transmissible, multi-antibiotic- and antiseptic-resistant variant of methicillin-resistant *Staphylococcus aureus*, sequence type 239 (TW). *J. Bacteriol.* **192**(3), 888–892 (2010).
- Laabei, M. *et al.* Predicting the virulence of MRSA from its genome sequence. *Genome Res.* **24**, 839–849 (2014).
- Saiki, K., Mogi, T. & Anraku, Y. Heme O biosynthesis in *Escherichia coli*: the *cyoE* gene in the cytochrome *bo* operon encodes a protoheme IX farnesyltransferase. *Biochem. Biophys. Res. Commun.* **189**(3), 1491–1497 (1992).
- Saiki, K., Mogi, T., Hori, H., Tsubaki, M. & Anraku, Y. Identification of the functional domains in heme O synthase. Site directed mutagenesis studies on the *cyoE* gene of the cytochrome *bo* operon in *Escherichia coli*. *J. Biol. Chem.* **268**(36), 26927–26934 (1993).
- Hill, J. *et al.* Demonstration by FTIR that the *bo*-type ubiquinol oxidase of *Escherichia coli* contains a heme-copper binuclear center similar to that in cytochrome *c* oxidase and that proper assembly of the binuclear center requires the *cyoE* gene product. *Biochemistry*. **31**, 11435–11440 (1992).
- Nakamura, H., Saiki, K., Mogi, T. & Anraku, Y. Assignment and functional roles of the *cyoABCDE* gene products required for the *Escherichia coli bo*-type quinol oxidase. *J. Biochem.* **122**, 415–421 (1997).
- Fey, P. D. *et al.* A genetic resource for rapid and comprehensive screening of nonessential *Staphylococcus aureus* genes. *mBio*. **4**(1), <https://doi.org/10.1128/mBio.00537-12> (2013).
- Tsuchiya, S. *et al.* Establishment and characterization of a human acute monocytic leukemia cell line (THP-1). *Int. J. Cancer*. **26**(2), 171–176 (1980).
- Kennedy, M. C., Emptage, M. H., Dreyer, J. L. & Beinert, H. The role of iron in the activation-inactivation of aconitase. *J. Biol. Chem.* **258**, 11098–11105 (1983).
- Biasini, M. *et al.* SWISS-MODEL: modelling protein tertiary and quaternary structure using evolutionary information. *Nucleic Acids Research* **42**(W1), W252–W258 (2014).
- Omasits, U., Ahrens, C. H., Müller, S. & Wollscheid, B. Protter: interactive protein feature visualization and integration with experimental proteomic data. *Bioinformatics*. **30**(6), 884–6 (2014).
- Recsei, P. *et al.* Regulation of exoprotein gene expression in *Staphylococcus aureus* by *agr*. *Mol. Gen. Genet.* **202**(1), 58–61 (1986).
- Janzon, L. & Avidson, Staffan. The role of the delta-lysine gene (*hld*) in the regulation of virulence genes by the accessory gene regulator (*agr*) in *Staphylococcus aureus*. *EMBO J.* **9**(5), 1391–1399 (1990).
- Arvidson, S. & Tegmark, K. Regulation of virulence determinants in *Staphylococcus aureus*. *Int. J. Med. Microbiol.* **291**, 159–170 (2001).
- Coleman, G. A comparison of the patterns of extracellular proteins produced by the high alpha-toxin-secreting organism *Staphylococcus aureus* (Wood 46) during aerobic and anaerobic growth. *J. Gen. Microbiol.* **131**(2), 405–408 (1985).
- Somerville, G. A. & Proctor, R. A. Cultivation conditions and the diffusion of oxygen into culture media: the rationale for the flask-to-medium ratio in microbiology. *BMC Microbiol.* **13**(9), <https://doi.org/10.1186/1471-2180-13-9> (2013).
- Thet, N. T. *et al.* Visible, colorimetric dissemination between pathogenic strains of *Staphylococcus aureus* and *Pseudomonas aeruginosa* using fluorescent dye containing lipid vesicles. *Biosens. Bioelectron.* **41**, 538–543 (2013).

Acknowledgements

E.S. was supported by funding from the BBSRC SWBio DTP. Thanks to Jean van den Elsen for his helpful advice. G.A.S. was supported by funds provided through the Hatch Act to the University of Nebraska Institute of Agriculture and Natural resources and by funds provided by Zoetis.

Author Contributions

E.S., M.L. and S.G. performed the experiments described here. E.S. and R.C.M. prepared the figures. E.S., G.A.S. and R.C.M. wrote the manuscript.

Additional Information

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017